

# Stratification and Sample Selection for Multicrop Experiments

by **M. M. Hixson**  
**B. J. Davis**  
**M. E. Bauer**

Principal Investigator  
**D. A. Landgrebe**

**November 1978**

Prepared for  
**National Aeronautics and Space Administration**

Johnson Space Center  
Earth Observation Division  
Houston, Texas 77058  
Contract No. NAS9-15466  
Technical Monitor: J. D. Erickson/SF3

Submitted by  
**Laboratory for Applications of Remote Sensing**  
**Purdue University**  
**West Lafayette, Indiana 47907**

# Stratification and Sample Selection for Multicrop Experiments

by M. M. Hixson  
B. J. Davis  
M. E. Bauer

(E79-10-190): STRATIFICATION AND SAMPLE  
SELECTION FOR MULTICROP EXPERIMENTS (Purdue  
Univ.) 53 p HC A04/MF A01 CSCL 02C

N79-24410

Unclas

G3/43 00190

Principal Investigator

**D. A. Landgrebe**

**November 1978**



Prepared for

**National Aeronautics and Space Administration**

Johnson Space Center  
Earth Observation Division  
Houston, Texas 77058

Contract No. NAS9-15466

Technical Monitor: J. D. Erickson/SF3

Submitted by

**Laboratory for Applications of Remote Sensing  
Purdue University**

**West Lafayette, Indiana 47907**

## STAR INFORMATION FORM

1 Report No. 112278	2. Government Accession No.	3. Recipient's Catalog No.	
4 Title and Subtitle Stratification and Sample Selection for Multicrop Experiments		5. Report Date 11/22/78	6. Performing Organization Code
		8. Performing Organization Report No LARS Contract Report 112278	
7. Author(s) Marilyn M. Hixson, Barbara J. Davis and Marvin E. Bauer		10. Work Unit No.	
9. Performing Organization Name and Address Laboratory for Applications of Remote Sensing Purdue University 1220 Potter Drive West Lafayette, Indiana 47906		11. Contract or Grant No. NAS9-15466	
		13. Type of Report and Period Covered	
12. Sponsoring Agency Name and Address J.D. Erickson/SF3 NASA/Johnson Space Center Houston, Texas 77058		14. Sponsoring Agency Code	
		15. Supplementary Notes	
16. Abstract <p>The purpose of this effort was to identify the locations of the sample segments for the 1978-79 Multicrop experiments to support:</p> <ul style="list-style-type: none"> <li>-Development and evaluation of procedures for using LACIE and other technologies for the classification of corn and soybeans.</li> <li>-Identification of factors likely to affect classification performance.</li> <li>-Evaluation of problems encountered and techniques which are applicable to the crop estimation problem in foreign countries as well.</li> </ul> <p>In order to meet these requirements, two types of samples were selected. Low density segments were distributed throughout corn and soybean producing areas to sample all variations of conditions which could affect classification accuracy and to more completely represent conditions which might be found in other countries. High density segments were selected in smaller areas to support the investigation of training, classification, and area estimation procedures on a smaller scale for possible use in future Multicrop experiments.</p> <p>In this report, the data set and methods employed in the stratification are discussed. Rationale, methods, and results for both the low and high density segments are discussed.</p>			
17 Key Words (Suggested by Author(s)) Stratification, Sampling, Landsat, Crop Inventory, corn and soybeans		18. Distribution Statement	
19 Security Classif (of this report) UNCLASSIFIED	20 Security Classif (of this page) UNCLASSIFIED	21. No. of Pages	22 Price*

## List of Tables

1. Crops included in the USDA/SRS county estimates program.	5
2. Determination of strata according to the normalized production of corn and soybeans and average farm size:	9
3. Counties assigned to Stratum 1.	18
4. Counties assigned to Stratum 2.	19
5. Counties assigned to Stratum 3.	20
6. Counties assigned to Stratum 4.	21
7. Counties assigned to Stratum 5.	23
8. Counties assigned to Stratum 6.	24
9. Counties assigned to Stratum 7.	25
10. Counties assigned to Stratum 8.	26
11. Locations of the low density sample segments by latitude and longitude coordinates of the segment centers.	30
12. Allocation of sample segments to counties in each of the four high density test sites.	40
13. Locations of the high density sample segments by latitude and longitude coordinates of the segment centers.	42
A-1. Commodity Codes.	46

## List of Figures

1. Record layout of county estimates data.	4
2. Schematic diagram illustrating the determination of strata for Multicrop experiments based on normalized production of corn and soybeans and average farm size.	8
3. Locations of counties assigned to Stratum 1, small farms, low production of corn and soybeans.	
4. Locations of counties assigned to Stratum 2, small farms, medium production of corn and soybeans.	11
5. Locations of counties assigned to Stratum 3, small farms, high production of corn and soybeans.	12
6. Locations of counties assigned to Stratum 4, large farms, low production of corn and soybeans.	13
7. Locations of counties assigned to Stratum 5, large farms, low production of corn, medium production of soybeans.	14
8. Locations of counties assigned to Stratum 6, large farms, medium production of corn and soybeans.	15
9. Locations of counties assigned to Stratum 7, large farms, high production of corn and soybeans.	16
10. Locations of counties assigned to Stratum 8, large farms, highest production of corn and soybeans.	17
11. Locations of counties in all eight strata receiving low density sample segments.	29
12. Locations of high density test sites.	37

## STRATIFICATION AND SAMPLE SELECTION FOR MULTICROP EXPERIMENTS

### 1. Introduction.

In February 1978, LARS was asked to participate in the stratification and sampling tasks for the transition year experiments. The project was supported by personnel and funds from two tasks of NASA Contract NAS9-15466: "Application of Statistical Pattern Recognition to Image Interpretation" and "Application and Evaluation of Landsat Training, Classification, and Area Estimation Procedures for Crop Inventory."

The purpose of this effort was to identify the locations of the sample segments for the 1978-79 Multicrop experiments to support:

- Development and evaluation of procedures for using LACIE and other technologies for the classification of corn and soybeans.
- Identification of factors likely to affect classification performance.
- Evaluation of problems encountered and techniques which are applicable to the crop estimation problem in foreign countries as well.

In order to meet these requirements, two types of samples were selected. Low density segments were distributed throughout corn and soybean producing areas to sample all variations of conditions which could affect classification accuracy and to more completely represent conditions which might be found in other countries. High density segments were selected in smaller areas to support the investigation of training, classification, and area estimation procedures on a smaller scale for possible use in future Multicrop experiments.

In this report, the data set and methods employed in the stratification are discussed. Rationale, methods, and results for both the low and high density segments are discussed.

## 2. Objectives

In order to support the corn and soybean experiments, two types of segments were selected: low density segments and high density segments. Different issues can be addressed using each type of segment.

The low density segments were selected to cover a wide range of conditions under which areas will have to be classified in larger Multicrop efforts to allow possible problems to be examined (e.g., in algorithms, systems, data acquisition). The low density samples were located in 14 states in the U.S. corn and soybean producing areas. This region was divided into eight strata according to the level of county production of corn and soybeans and average farm size. Twenty segments per stratum were selected. The distribution of these segments permits the calculation of variability within a stratum to predict the variability of aggregated estimates of corn and soybeans in the U.S. and to determine the optimum allocation of samples for making such estimates. The allocation of these samples was not designed for, and thus does not support, making aggregated estimates.

The high density samples are located in four test sites in high production areas of the U.S. Corn Belt. Twenty segments were selected from each test site which is approximately ten counties in size. The increased density of samples permits estimation of the local variability in high production areas. These samples support the investigation of training, classification, and area estimation procedures on a smaller scale for possible use in future Multicrop experiments. Other area estimation procedures such as regression estimation can be evaluated and county level estimates can be assessed.

## 3. Data Set Description

The data used in this study were acquired by the Statistical Reporting Service of the U.S. Department of Agriculture (USDA/SRS). Two types of data were available: the USDA/SRS county estimates for 1972-76 and the 1974 agriculture census data. The data were supplied by NASA/Johnson Space Center (NASA/JSC).

The SRS dual county estimates program data for 1972-76 were available. Under the Federal program, county estimates are prepared for specified crops, states, and counties. These estimates include the major crops produced in most states. Some of the state statistical offices prepare county estimates for a few crops not required under the Federal program in cooperation with their respective state governments, but these estimates were not available on tape.

Variables which were included in the county estimates data set were: state, crop reporting district, county, year data was punched, crop year, commodity code, acres planted, acres harvested, yield per harvested acre, and production (Figure 1). Counties from the entire U.S. were represented. The commodities for which information was available are listed in Table 1. Some problems encountered with this data set are discussed in the appendix.

The 1974 agriculture census data were supplied for 14 states in the U.S. corn and soybean producing regions. These data included: number of acres in each county, average farm size by county, and the land in farms for each county.

#### 4. Stratification

The first step in selection of sample segments was the stratification of the area to be studied. The variables used in the stratification, the rationale and methods employed, and the results of the stratification will be discussed in this section.

##### Variables Used in Stratification.

The variables available were those contained in the USDA/SRS county estimates program (Figure 1) and the selected variables from the 1974 agriculture census which were supplied by NASA/JSC. The variables which were considered for use were: acres planted, acres harvested, yield, and production for the crops listed in Table 1; acres in a county; percent agricultural area (land in farms) in a county; and average farm size by county. From these variables, the



CARD NO.	SURVEY CODE	ID					DATA																					
		STATE	DISTRICT	COUNTY	YEAR PUNCHED	CROP YEAR	COMMODITY CODE	(1) ACRES PLANTED ALL PURPOSES	(2) ACRES PLANTED FOR HARV. OR NET ACRES SEEDED OR ACRES ABANDONED	(3) ACRES HARVESTED	(4)	(5) YIELD PER HARVESTED ACRE	(6) PRODUCTION	(7) PRODUCTION (COTTON LBS OF LINT)														
1	2	5	6	7	8	9	10	12	13	14	15	16	17	26	27	34	35	42	43	50	51	54	55	58	59	68	69	78

Figure 1. Record layout of county estimates data.



number of agricultural acres in a county was computed by multiplying the percent agricultural area by the county acreage. Normalized production of a crop for a county was computed by dividing the five-year average production of that crop by the agricultural acres in the county.

In order to fulfill the objectives, the stratification was performed using three variables: normalized production of corn, normalized production of soybeans, and average farm size. The first two variables were selected to make strata which are homogeneous with respect to the relative importance of corn and soybeans in the agricultural scene. The average farm size was selected to represent problems which might be encountered in Landsat data classifications with different field sizes.

#### Methods of Stratification.

The rationale for the stratification method was based upon the objective of creating eight strata in the United States corn and soybean producing regions which were relatively homogeneous with respect to the relative importance of corn and soybeans in the agricultural scene and the average farm (or field) size. These strata, then, represent several conditions under which Landsat data will have to be classified in Multicrop studies. Samples selected from these strata will be representative of conditions found throughout the corn and soybean producing regions.

The first step in the stratification was a reduction of the data set size. Only the 14 states for which the agriculture census data were supplied were considered. Counties with neither corn nor soybeans were omitted.

The joint distributions of normalized corn and soybean productions and average farm size were examined. The average farm size was represented in two groups: small farms (average size less than or equal to 190 acres) and large farms (size greater than 190 acres).

About one-third of the counties were in the small farms category and about two-thirds were in the large farms category. The division into these two groups was somewhat arbitrary although there was a break in the continuum of data at about 190 acres.

For each farm size, the normalized corn and soybean productions were displayed in deciles to look for broad clusters of data. The strata were determined by examining tables of the distributions of these variables. Three strata of small farm counties and five strata of large farm counties were selected to represent the two farm sizes approximately proportionally to the number of counties in them.

Counties which fell in the lower 10% of all counties in both corn and soybean production were omitted from consideration. Counties which fell outside the broad clusters of data were not included in any stratum. Thirteen counties satisfying all other selection criteria were outliers from the clusters and were not included. A schematic diagram (Figure 2 ) shows the methodology employed in the stratification. Table 2 gives the definitions of stratum boundaries.

#### Results of Stratification.

Eight strata covering 14 states in the U.S. corn and soybean producing region were determined. The counties in each of these strata are shown in Figures 3 to 10 and are listed in Tables 3 to 10.

The large farm, highest production stratum (stratum 8) is geographically located at the center of the Corn Belt. Strata 7, 6, and 4 are located around its perimeter outward according to decreased production. In these strata of large farms, corn and soybeans are of approximately equal importance.

Stratum 5 is located geographically apart from the other strata with large farms. This stratum, in which soybeans have a greater





Figure 2. Schematic diagram illustrating the determination of strata for Multicrop experiments based on normalized production of corn and soybeans and average farm size.

Table 2. Determination of strata according to the normalized production of corn and soybeans and average farm size.

Stratum Number	Average Farm Size	Normalized Production		No. of Counties
		Corn	Soybeans	
	(acres)	(deciles)	(deciles)	
1	<190	0-40	0-40	149
2	<190	40-60	30-70	109
3	<190	60-100	50-100	126
4	>190	0-40	0-30	192
5	>190	0-40	30-70	102
6	>190	40-60	30-70	126
7	>190	60-80	50-90	147
8	>190	80-100	70-100	213




Figure 3. Locations of counties assigned to Stratum 1, small farms, low production of corn and soybeans.




Figure 4. Locations of counties assigned to Stratum 2, small farms, medium production of corn and soybeans.








Figure 6. Locations of counties assigned to Stratum 4, large farms, low production of corn and soybeans.






Figure 8. Locations of counties assigned to Stratum 6, large farms, medium production of corn and soybeans.




Figure 9. Locations of counties assigned to Stratum 7, large farms, high production of corn and soybeans.




Figure 10. Locations of counties assigned to Stratum 8, large farms, highest production of corn and soybeans.

Table 3. Counties assigned to Stratum 1.

<u>Arkansas</u>	<u>Kentucky</u>	<u>Michigan</u>	<u>Missouri</u>
Benton	Knox	Alpena	St. Francois
Bradley	Laurel	Benzie	Webster
Calhoun	Letcher	Crawford	
Cleburne	Lewis	Gogebic	<u>Ohio</u>
Cleveland	McCreary	Leelanau	Athens
Columbia	Madison	Manistee	Belmont
Crawford	Magoffin	Oceana	Guernsey
Garland	Martin	Wexford	Jefferson
Grant	Menifee		Lake
Hot Springs	Montgomery	<u>Mississippi</u>	Lawrence
Howard	Morgan	Choctaw	Meigs
Johnson	Nicholas	Clarke	Monroe
Montgomery	Owen	Covington	
Pike	Owsley	Forrest	<u>Wisconsin</u>
Saline	Pendleton	Greene	Clark
Union	Perry	Harrison	Door
	Pike	Itawamba	Kewaunee
<u>Kentucky</u>	Powell	Jackson	Marathon
Adair	Robertson	Jasper	Wood
Allen	Rockcastle	Jeff Davis	
Anderson	Rowan	Jones	
Bath	Scott	Leake	
Boone	Trimble	Lincoln	
Boyd	Washington	Neshoba	
Bracken	Wayne	Newton	
Breathitt	Whitley	Perry	
Bullitt	Wolfe	Pike	
Campbell	Woodford	Pontotoc	
Carroll		Scott	
Carter	<u>Louisiana</u>	Simpson	
Clay	Bienville	Smith	
Cumberland	Grant	Stone	
Elliott	Jackson	Tippah	
Estill	Lafayette	Tishomingo	
Fayette	La Salle	Union	
Floyd	Lincoln	Walthall	
Franklin	Livingston	Wayne	
Gallatin	Sabine	Winston	
Garrard	St. Helena		
Grant	St. Martin	<u>Missouri</u>	
Greenup	St. Tammany	Barry	
Harlan	Tangipahoa	Christian	
Harrison	Union	Cole	
Jackson	Vernon	Greene	
Jessamine	Washington	Jefferson	
Johnson	Webster	Lawrence	
Kenton	Winn	Newton	
Knott			

Table 4. Counties assigned to Stratum 2.

<u>Illinois</u>	<u>Kentucky</u>	<u>Ohio</u>
Jefferson	Monroe	Cuyahoga
Williams	Muhlenberg	Gallia
	Nelson	Geauga
<u>Indiana</u>	Pulaski	Hamilton
Brown	Russell	Hocking
Clark	Shelby	Jackson
Crawford	Spencer	Lorain
Dearborn	Taylor	Muskingum
Floyd	Warren	Ottawa
Harrison		Perry
Jefferson	<u>Michigan</u>	Scioto
Lawrence	Arenac	Summit
Monroe	Bay	Trumbull
Ohio	Gladwin	Tuscarawas
Perry	Grand Traverse	Washington
Switzerland	Kent	<u>Wisconsin</u>
	Mason	Brown
<u>Kentucky</u>	Montcalm	Calumet
Ballard	Muskegon	Manitowoc
Barren	Newaygo	Milwaukee
Boyle	Oakland	Oconto
Breckinridge	Ottawa	Outagami
Caldwell	Saginaw	Ozaukee
Calloway	St. Clair	Polk
Carlisle	Sanilac	Shawano
Casey	Van Buren	Sheboyga
Clark	Wayne	Vernon
Clinton		Waupaca
Edmonson	<u>Minnesota</u>	
Graves	Anoka	
Grayson	Benton	
Green	Isanti	
Hancock	Mille Lacs	
Hardin	Ramsey	
Hart		
Henry	<u>Missouri</u>	
Jefferson	Franklin	
Larue	Jackson	
Lincoln	St. Louis	
Logan		
Lyon	<u>Ohio</u>	
McCracken	Adams	
Marion	Ashtabula	
Marshall	Brown	
Mason	Carroll	
Meade	Clermont	
Mercer		
Metcalf		



Table 5. Counties assigned to Stratum 3.

<u>Indiana</u>	<u>Kentucky</u>	<u>Ohio</u>
Calhoun	Davies	Burler
Cook	Fleming	Columbiana
Madison	McLean	Crawford
Massac	Ohio	Darke
Adams	Simpson	Defiance
Allen		Erie
Blackford	<u>Michigan</u>	Fairfield
Daviess	Allegan	Fulton
De Kalb	Barry	Henry
Delaware	Berrien	Highland
Dubois	Branch	Holmes
Elkhart	Cass	Knox
Franklin	Clinton	Licking
Grant	Eaton	Logan
Hamilton	Genesee	Lucas
Hancock	Gratiot	Mahoning
Hendricks	Hillsdale	Mercer
Henry	Ingham	Miami
Howard	Ionia	Montgomery
Huntington	Isabella	Morrow
Jackson	Kalamazoo	Portage
Jay	Lapeer	Preble
Jennings	Lenawee	Putnam
Johnson	Livingston	Richland
Kosciusko	Macomb	Sandusky
Lagrange	Midland	Shelby
Madison	Monroe	Stark
Marion	Shiawassee	Warren
Marshall	Tuscola	Wayne
Miami	Washtenaw	Williams
Morgan		
Noble	<u>Minnesota</u>	
Orange	Carver	
Owen	Chisago	
Pike	Hennepin	
Randolph	McLeod	
Ripley	Rice	
St. Joseph	Scott	
Scott	Steele	
Spencer	Washington	
Steuben	Wright	
Washington		
Wayne	<u>Ohio</u>	
Wells	Allen	
Whitley	Ashland	
	Auglaize	
<u>Iowa</u>		
Bremer		

Table 6. Counties assigned to Stratum 4.

<u>Arkansas</u>	<u>Michigan</u>	<u>Mississippi</u>
Clark	Alcona	Hancock
Dallas	Alger	Hinds
Faulkner	Antrim	Jefferson
Franklin	Cheboygan	Kemper
Hempstead	Delta	Lafayette
Izard	Dickinson	Lamar
Lafayette	Emmet	Lauderdale
Little Rock	Houghton	Lawrence
Logan	Kalkaska	Madison
Nevada	Lake	Marion
Ouachita	Luce	Marshall
Perry	Mackinac	Montgomery
Pope	Marquette	Oktibbeha
Scott	Menominee	Pearl River
Sebastian	Montmorenci	Rankin
Sevier	Osceola	Wilkinson
Yell	Oscoda	Yalobush
	Otsego	
<u>Kentucky</u>	Presque	<u>Missouri</u>
Lawrence	Schoolcraft	Benton
		Camden
<u>Louisiana</u>	<u>Minnesota</u>	Carter
Ascension	Aitkin	Cedar
Assumption	Becker	Crawford
Beauregard	Beltrami	Dade
Bossier	Carlton	Dallas
Caddo	Cass	Dent
Calcasieu	Clay	Douglas
Caldwell	Clearwater	Hickory
Cameron	Crow Wing	Howell
Claiborne	Hubbard	Iron
De Soto	Itasca	LaCledde
East Baton Rouge	Mahnomen	Madison
East Feliciana	Norman	Maries
Iberia	Pennington	Miller
Iberville	Pine	Morgan
LaFourche	Polk	Oregon
Natchitoches	Red Lake	Osage
Plaquemines	Wilkin	Phelps
Red River		Polk
St. James	<u>Mississippi</u>	Pulaski
St. Mary	Amite	Reynolds
Terrebon	Attala	Ripley
Vermilion	Carroll	Shannon
West Baton Rouge	Claiborne	Stone
West Feliciana	Copiah	Texas
	De Soto	Washington
	Franklin	Wayne
		Wright

Table 6. (con't.)

<u>Nebraska</u>	<u>South Dakota</u>
Banner	Potter
Blaine	Spink
Box Butte	Sully
Cherry	Todd
Cheyenne	Tripp
Dawes	Walworth
Deuel	
Garden	<u>Wisconsin</u>
Garfield	
Keith	Florence
Keya Paha	Forest
Kimball	Iron
Logan	Langlade
Loup	Lincoln
McPherson	Rusk
Morrill	Sawyer
Rock	Taylor
Sheridan	Washburn
Sioux	
Thomas	
<u>Ohio</u>	
Morgan	
Noble	
<u>South Dakota</u>	
Aurora	
Beadle	
Brown	
Brule	
Buffalo	
Butte	
Campbell	
Clark	
Codington	
Custer	
Day	
Edmunds	
Fall River	
Faulk	
Gregory	
Hand	
Hughes	
Hyde	
Jerauld	
Lyman	
McPherson	
Marshall	

Table 7. Counties assigned to Stratum 5.

<u>Arkansas</u>	<u>Michigan</u>	<u>Missouri</u>
Arkansas	Charlevoir	New Madrid
Ashley	Iosco	Pemiscot
Chicot		Putnam
Clay	<u>Minnesota</u>	St. Clair
Conway		Vernon
Craighead	Kanabec	
Crittenden	Otter	<u>Nebraska</u>
Cross	Traverse	Boyd
Desha	Wadena	Lancaster
Drew		
Greene	<u>Mississippi</u>	<u>Ohio</u>
Independence	Adams	Vinton
Jackson	Benton	
Jefferson	Bolivar	
Lawrence	Calhoun	<u>South Dakota</u>
Lee	Chickasaw	Charles Mix
Lincoln	Clay	Grant
Lonoke	Coahoma	Miner
Miller	Grenada	Roberts
Mississippi	Holmes	Sanborn
Monroe	Humphrey	
Phillips	Issaquen	<u>Wisconsin</u>
Poinsett	Lee	Barron
Prairie	Leflore	Burnett
Pulaski	Lowndes	Marinette
Randolph	Monroe	
St. Francis	Noxubee	
White	Panola	
Woodruff	Prentiss	
	Quitman	
<u>Louisiana</u>	Sharkey	
Adadia	Sunflower	
Allen	Tallahatchie	
Catahoula	Tate	
Concordia	Tunica	
East Carroll	Warren	
Evangeline	Washington	
Franklin	Webster	
Jefferson	Yazoo	
Madison		
Morehouse	<u>Missouri</u>	
Ouachita	Barton	
Pointe Coupee	Bollinger	
Rapides	Butler	
Richland	Dunklin	
St. John	Gasconade	
Tensas	Henry	
	Jasper	

Table 8. Counties assigned to Stratum 6.

<u>Illinois</u>	<u>Missouri</u>	<u>Nebraska</u>	<u>Wisconsin</u>
Alexander	Bates	Frontier	Pierce
Franklin	Boone	Furnas	Portage
Hardin	Caldwell	Hitchcock	Richland
Jackson	Callaway	Holt	St. Croix
Johnson	Cape Girardeau	Jefferson	Trempealeau
Monroe	Carroll	Johnson	Waushara
Perry	Cass	Knox	
Pope	Chariton	Lincoln	
Pulaski	Clay	Nuckolls	
Randolph	Clinton	Pawnee	
Union	Cooper	Perkins	
	Daviess	Red Willow	
<u>Iowa</u>	De Kalb	Sherman	
Clarke	Gentry	Webster	
Decatur	Grundy	Wheeler	
	Harrison		
	Howard	<u>Ohio</u>	
	Johnson	Harrison	
	Knox	Pike	
<u>Kentucky</u>	Lewis		
Bourbon	Linn		
Butler	Livingston	<u>South Dakota</u>	
Christian	Macon	Bon Homme	
Crittenden	Marion	Brookings	
Livingston	Mercer	Davison	
Oldham	Moniteau	Deuel	
Trigg	Monroe	Douglas	
	Montgomery	Hamlin	
<u>Michigan</u>	Perry	Hanson	
Clarey	Pettis	Hutchins	
Mecosta	Pike	Kingsburg	
Missaukee	Platte	Lake	
Ogemaw	Ralls	McCook	
	Randolph		
<u>Minnesota</u>	Ray	<u>Wisconsin</u>	
Big Stone	Ste. Genevieve	Adams	
Douglas	Schuyler	Buffalo	
Grant	Scotland	Chippewa	
Morrison	Scott	Crawford	
Pope	Shelby	Dunn	
Sherburne	Stoddard	Eau Clair	
Stearns	Sullivan	Jackson	
Todd	Warren	Juneau	
	Worth	La Crosse	
<u>Missouri</u>	<u>Nebraska</u>	Marquette	
Adair	Brown	Monroe	
Audrain	Custer	Pepin	

Table 9. Counties assigned to Stratum 7.

<u>Illinois</u>	<u>Iowa</u>	<u>Missouri</u>	<u>Ohio</u>
Adams	Van Buren	Lafayette	Paulding
Bond	Wapello	Lincoln	Pickaway
Brown	Warren	Nodaway	Ross
Clark	Wayne	St. Charles	Seneca
Clay	Winneshiek	Saline	Union
Clinton			Wyandot
Crawford	<u>Kentucky</u>	<u>Nebraska</u>	<u>South Dakota</u>
Cumberland	Henderson	Antelope	Clay
Edwards	Hickman	Boone	Lincoln
Effingham	Hopkins	Buffalo	Minnehaha
Fayette	Todd	Butler	Moody
Hamilton	Webster	Cass	Turner
Jasper		Cedar	Union
Jersey	<u>Michigan</u>	Chase	Yankton
Jo Daviess	Calhoun	Colfax	
Lake	Huron	Cuming	<u>Wisconsin</u>
Marion	Jackson	Dakota	Columbia
Pike	St. Joseph	Dawson	Franklin
Richland		Dixon	Grant
St. Clair	<u>Minnesota</u>	Franklin	Green
Saline	Chippewa	Gosper	Green Lake
Schuyler	Dakota	Greeley	Iowa
Washington	Fillmore	Harlan	Lafayette
Wayne	Goodhue	Howard	Sauk
White	Houston	Madison	Walworth
	Kandiyohi	Nance	
<u>Indiana</u>	Lac Qui	Nemaha	
Greene	Lincoln	Otoe	
Martin	Lyon	Pierce	
Warrich	Meeker	Richards	
	Murray	Saline	
<u>Iowa</u>	Olmsted	Saunders	
Adair	Pipestone	Seward	
Adams	Redwood	Stanton	
Allamakee	Stevens	Thayer	
Appanoose	Swift	Thurston	
Davis	Wabasha	Valley	
Guthrie	Winona	Washington	
Howard	Yellow Medicine	Wayne	
Jackson		<u>Ohio</u>	
Lucas	<u>Missouri</u>	Coshocton	
Madison	Andrew	Delaware	
Marion	Atchison	Franklin	
Monroe	Buchanan	Hancock	
Page	Clark	Hardin	
Ringgold	Holt	Huron	
Taylor		Marion	
Union			

Table 10. Counties assigned to Stratum 8.

<u>Illinois</u>	<u>Illinois</u>	<u>Iowa</u>
Boone	Stark	Audubon
Bureau	Stephens	Benton
Carroll	Tazewell	Black Hawk
Cass	Vermilion	Boone
Champaign	Wabash	Buchanan
Christian	Whiteside	Buena Vista
Coles	Will	Butler
De Kalb	Winnebago	Calhoun
De Witt	Woodford	Carroll
Douglas		Cass
Du Page	<u>Indiana</u>	Cedar
Edgar	Bartholomew	Cerro Gordo
Ford	Benton	Cherokee
Fulton	Boone	Chickasaw
Gallatin	Carroll	Clay
Greene	Cass	Clayton
Grundy	Clay	Clinton
Hancock	Clinton	Crawford
Henderson	Decatur	Dallas
Henry	Fayette	Delaware
Iroquois	Fountain	Dès Moines
Kane	Fulton	Dickinson
Kankakee	Gibson	Dubuque
Kendall	Jasper	Emmet
Knox	Knox	Fayette
La Salle	Lake	Floyd
Lawrence	La Porte	Franklin
Lee	Montgomery	Fremont
Livingston	Newton	Greene
Logan	Parke	Grundy
McDonough	Porter	Hamilton
McHenry	Posey	Hancock
McLean	Pulaski	Hardin
Macon	Putnam	Harrison
Macoupin	Rush	Henry
Marshall	Shelby	Humboldt
Mason	Starke	Ida
Menard	Sullivan	Iowa
Mercer	Tippecanoe	Jasper
Montgomery	Tipton	Jefferson
Morgan	Union	Johnson
Moultrie	Vanderburg	Jones
Ogle	Vermillion	Keokuk
Peoria	Vigo	Kossuth
Piatt	Wabash	Lee
Putnam	Warren	Linn
Rock Island	White	Louisa
Sangamon		Lyon
Scott		Mahaska
Shelby		Marshall

Table 10. (con't.)

<u>Iowa</u>	<u>Nebraska</u>
Mills	Adams
Mitchell	Burt
Monona	Clay
Montgomery	Dodge
Muscatine	Douglas
O'Brien	Fillmore
Osceola	Hall
Palo Alto	Hamilton
Plymouth	Kearney
Pocahontas	Merrick
Polk	Phelps
Pottawattamie	Platte
Poseshiek	Polk
Sac	Sarpy
Scott	York
Shelby	
Sioux	<u>Ohio</u>
Story	Champaign
Tama	Clark
Washington	Clinton
Webster	Fayette
Winnebago	Greene
Woodbury	Madison
Worth	Van Wert
Wright	Wood
	<u>Wisconsin</u>
<u>Kentucky</u>	Rock
Union	
<u>Minnesota</u>	
Blue Earth	
Brown	
Cottonwood	
Dodge	
Faribault	
Freeborn	
Jackson	
Le Sueur	
Martin	
Mower	
Nicollet	
Nobles	
Renville	
Rock	
Sibley	
Waseca	
Watonwan	



importance than corn, is located in the Mississippi River Valley where the climate and soils are more suited to soybeans than to corn.

Stratum 3, the small farm stratum with the greatest production of corn and soybeans, is located primarily in eastern Indiana and western Ohio where the cropland is productive, but the terrain is rolling. The lesser production small farm strata (strata 1 and 2) are centered about this area on the outskirts of stratum 3.

In summary, looking at the geographic location of the strata, the system appears to be logical and the various strata seem to represent different conditions. This result is supportive not only of the variables and the methodology employed in the stratification, but also of the validity of the data sets employed.

#### 5. Low Density Segments

##### Sample Allocation.

The low density segments were selected to sample the variability present in corn and soybean producing regions of the United States. The sample was designed to represent differences in climate, topography, field size, variety, and management practices. In order to achieve as diverse a representation as possible, an equal number of segments were allocated to each of the strata. This allocation scheme emphasizes representation of variability rather than sampling in a manner suitable for aggregation purposes.

Twenty 5 x 6 nautical mile segments were allocated to each stratum. The counties to receive sample segments were determined using a random selection procedure without replacement. Thus, all counties in a stratum had an equal probability of receiving a sample and no county could contain more than one segment. Locations of counties receiving sample segments are illustrated in Figure 11. Latitude and longitude coordinates of the sample segment centers can be found in Table 11.




Figure 11. Locations of counties in all eight strata receiving low density sample segments.

Table 11. Locations of the low density sample segments by latitude and longitude coordinates of the segment centers.

Stratum	State	County	Latitude/Longitude
1	Arkansas	Clebuine	35.480/91.970
	Kentucky	Bath	38.240/83.807
		Boone	38.815/84.675
		Boyd	38.385/82.658
		Carter	38.312/83.170
		Clay	37.237/83.833
		Greenup	38.643/82.933
		Lewis	38.570/83.545
		Nicholas	38.327/84.060
		Powell	37.823/83.817
		Scott	38.223/84.636
		Washington	37.677/85.081
		Michigan	Leelanau
	Mississippi	Jones	31.594/89.204
		Ponototoc	34.300/89.082
		Smith	32.012/89.436
	Missouri	Greene	37.164/93.470
		St. Francis	37.880/90.540
	Ohio	Belmont	40.112/81.000
	Wisconsin	Wood	44.500/90.000
2	Illinois	Jefferson	38.340/89.101
	Indiana	Switzerland	38.858/85.033
	Kentucky	Ballard	37.117/88.963
		Barren	37.065/85.885
		Graves	36.706/88.695
		Grayson	37.413/86.590
		Lyon	37.058/87.960
		McCracken	37.030/88.735
		Monroe	36.715/85.710
	Michigan	Arenac	44.086/83.685
		Kent	43.271/85.763
	Missouri	Jackson	38.908/94.263
	Ohio	Adams	38.960/83.470
		Ashtabula	41.696/80.818
		Gallia	38.876/82.317
		Jackson	38.930/82.573
		Sciota	38.795/82.829
		Washington	39.456/81.665

Table 11. (con't)

Stratum	State	County	Latitude/Longitude	
3	Wisconsin	Polk	45.283/92.283	
		Vernon	43.617/90.900	
	Illinois	Calhoun	38.920/90.575	
		Indiana	Daviess	38.791/87.102
			Hamilton	40.127/86.070
			Jennings	39.040/85.563
			Johnson	39.415/86.245
			Scott	38.693/85.725
			Spencer	37.975/87.145
			Washington	38.624/86.080
			Whitley	41.127/85.667
		Kentucky	Fleming	38.423/83.750
	Daviess		37.660/87.125	
	Michigan	Ingham	42.665/84.278	
	Minnesota	Carver	44.766/93.800	
		Scott	44.633/93.383	
		Wright	45.150/93.900	
Ohio	Defiance	41.372/84.550		
	Fulton	41.541/84.288		
	Logan	40.463/83.612		
	Portage	41.284/81.230		
4	Wisconsin	Dane	42.922/89.385	
	Louisiana	Cameron	29.950/93.080	
		De Soto	32.110/93.790	
		East Baton Rouge	30.670/91.095	
		Iberville	30.141/91.155	
		Red River	32.173/93.360	
		West Feliciana	30.805/91.315	
	Minnesota	Polk	47.816/96.683	
	Mississippi	Carroll	33.345/89.813	
		Copiah	31.761/90.611	
		Hinds	32.348/90.615	
		Yalobusha	34.140/89.635	
	Missouri	Dent	37.628/91.600	
		Laclede	37.669/92.595	
		Polk	37.667/93.351	
		Texas	37.283/92.000	
Nebraska	Box Butte	42.166/103.233		
	Keith	41.168/101.866		
	Kimball	41.307/103.650		

Table 11. (con't)

Stratum	State	County	Latitude/Longitude
5	South Dakota	Aurora	43.750/98.483
		Hyde	44.466/99.450
	Arkansas	Conway	35.190/92.790
		Greene	36.192/90.710
		Jefferson	34.354/91.882
		Lonoke	34.772/92.003
		Prairie	34.762/91.615
		White	35.208/91.580
	Louisiana	Allen	30.490/92.815
		Madison	32.282/91.501
		Morehouse	32.910/91.630
	Minnesota	Traverse	45.819/96.451
		Wadena	46.439/94.897
	Mississippi	Benton	34.937/89.295
		Calhoun	33.932/89.183
Humphreys		33.305/90.365	
Noxobe		33.191/88.543	
Sharkey		32.750/90.880	
Tunica		34.570/90.305	
Yazoo		32.765/90.143	
South Dakota	Roberts	45.725/96.950	
	Sanborn	43.996/97.878	
6	Illinois	Pope	37.335/88.605
	Iowa	Decatur	40.631/94.014
	Kentucky	Crittendon	37.245/88.150
	Michigan	Mecosta	43.681/85.206
	Missouri	Adair	40.250/92.500
		Boone	39.215/92.183
		Callaway	38.962/92.035
		Clay	39.410/94.276
		Cooper	38.745/92.870
		Gentry	40.325/94.430
		Grundy	40.171/93.381
Lewis		40.005/91.670	
Mercer		40.338/93.383	
Platte	39.484/94.795		
South Dakota	Brookings	44.304/96.890	
	Deuel	44.963/96.570	
	Douglas	43.333/98.179	

Table 11. (con't)

Stratum	State	County	Latitude/Longitude
7	Wisconsin	Crawford	43.127/91.034
		Eau Claire	44.735/91.255
		Trempeal	44.387/91.360
	Illinois	Hamilton	38.035/88.495
		Pike	39.665/91.210
		Richland	38.695/88.135
		St. Claire	38.589/89.865
	Iowa	Lucas	41.050/93.489
		Madison	41.466/94.021
		Taylor	40.792/94.806
		Wapello	40.959/92.300
		Warren	41.384/93.489
	Kentucky	Hickman	36.698/88.944
	Michigan	Jackson	42.336/84.425
	Minnesota	Goodhue	44.453/92.875
Missouri	Atchison	40.310/95.214	
	Clark	40.360/91.520	
	Lincoln	39.080/91.130	
Nebraska	Antelope	42.367/98.180	
	Dawson	40.908/99.955	
	Dixon	42.333/96.916	
Ohio	Delaware	40.212/82.826	
	Wyandot	40.880/83.352	
8	Illinois	Boone	42.178/88.809
		Douglas	39.749/88.055
		LaSalle	41.428/89.083
		Logan	40.259/89.221
		McLeon	40.675/88.824
		Moultrie	39.755/88.703
	Indiana	Carroll	40.712/86.593
		Gibson	38.288/87.352
		Lake	41.294/87.345
		Montgomery	40.211/86.854
		Vermilion	39.622/87.498
	Iowa	Butler	42.717/92.674
		Chickasaw	43.131/92.395
		Floyd	43.134/92.805
		Jefferson	41.122/91.900
Kossuth		43.299/94.310	
O'Brien		43.035/95.399	

Table 11. (con't)

Stratum	State	County	Latitude/Longitude
	Minnesota	Cottonwood	44.016/95.133
		Freeborn	43.787/93.429
	Ohio	Clinton	39.377/83.602

### Segment Location.

Segment locations were selected using a modification of a computer program written for "Crop Inventory Using Full-Frame Classification", described in the final report of Contract NAS9-14970 (June, 1977). The design of the location procedure was based upon that used in LACIE. A grid was laid over each county with grid intersections five by six nautical miles apart. A random selection procedure was then used to select a grid intersection which determined the latitude and longitude coordinates of the center point of each segment.

Although only one segment was allocated to each county, several segments were selected to attain a high probability that at least one of them would be located in an agricultural area and would be accepted as a site. The number of sites to be located in each county was determined by the percent agricultural land in the county. The segment centers were randomly selected without replacement and the first segment located outside a nonagricultural area was to be used.

The ag/nonag delineation was conducted by NASA/JSC. Full-frame color composite Landsat imagery was used to delineate areas which were not agricultural. This was done on the basis of whether or not field patterns were apparent. Rangeland, forest, and urban areas were among the types of land uses which were delineated as nonag. Segment locations were compared with these boundaries and the segment was rejected if less than 5% of the segment fell into an agricultural area.

## 6. High Density Segments

### Test Site Selection.

The high density segments were designed for intensive study of the remote sensing technology required for corn and soybean inventories. In order to sample more corn and soybeans, test sites were located in the Corn Belt where production of both crops is high. Test sites were



placed across the Corn Belt to sample the varied climatic conditions, soil types, crop distributions, and field sizes which are present (Figure 12). Each test site was selected to be relatively homogeneous within (same stratum, similar soil types and farming practices) to support classification studies, particularly of multisegment training. Each of the sites contained about ten counties and was approximately the size of a crop reporting district.

Test Site 1 is located in eastern Indiana which is an area of small farms. The other three test sites are located in large farm areas. Test Site 2 is comprised of counties in west central Indiana and east central Illinois. Test Site 3 is in north central Iowa and Test Site 4 is in west central Iowa.

Description of Test Sites 1 and 2. The climate across central Indiana and east central Illinois is continental with warm summers and cold winters. Normal mean temperature is  $-1.2^{\circ}\text{C}$  in January and  $31.1^{\circ}\text{C}$  in July. In this semihumid region of the U.S., the average annual precipitation is 950 to 1000 mm which does not limit crop production. Rainfall is greatest during the spring and early summer months with June typically receiving 107 to 118 mm of rain. Average precipitation in June is slightly excessive, adequate in July, and often inadequate in August for corn. The crops survive because of some moisture stored in the soil profile.

Test Site 1 is composed of two major soil associations. Soils of the northern two-thirds of this district (Allen, Wells, Adams, Blackford, Jay, and parts of Madison, Delaware, and Randolph counties) belong to the Blont-Pewano-Mortley soil association. These soils were formed on clayey glacial till and are nearly level and poorly to very poorly drained. The Brookston-Crosby-Miami-Parr association which predominates in the remainder of Test Site 1 was formed in thin loess (wind-blown materials) over loamy glacial till and is also poorly drained. These two soil associations are suited to intensive cropping but are subject to problems associated with wet soils unless adequate artificial drainage is provided. Typically, approximately 287,700 hectares of




Figure 12. Locations of high density test sites.

corn for grain; 245,300 hectares of soybeans; and 87,300 hectares of winter wheat are planted.

Test Site 2 includes dark-colored prairie soils and light-colored forest soils both of which were formed in loess-covered glacial till. Topography is generally gently rolling with short slopes and nearly level areas interrupted by depressions or potholes. The northern one-third of this district (Newton, Jasper, Kankakee, and northern Ford and Iroquois counties) has soils which are sandy and variable in subsoil development. These soils tend to be droughty, low in fertility, and require a high level of management for moderate yields. In Tippecanoe, Benton, Warren, southern Ford and Iroquois, and northern Vermilion and Champaign counties in the central portion of the district, the soils developed under prairie or mixed prairie and forest vegetation, are dark to moderately dark colored, and are generally imperfectly drained. Crop yields are moderately high to high with a high level of management. Dark-colored soils on nearly level to moderately sloping upland areas are typical in southern Vermilion and Champaign counties. These soils have high available moisture storage capacities and are very highly productive under a high level of management. Farmers in Test Site 2 typically plant 667,700 hectares of corn; 557,200 hectares of soybeans; and 39,200 hectares of winter wheat.

Description of Test Sites 3 and 4. The climate in western Iowa is continental, characterized by marked seasonal changes. Temperature fluctuations are extreme with winters being cold and summers warm. Thirty-year normal temperatures are  $-8.4^{\circ}\text{C}$  in January, the coldest month, and  $23.6^{\circ}$  in July, the warmest month. Annual precipitation is 762 mm with most of it occurring in the spring and early summer. Summer precipitation is variable from year to year with the largest amount (132 mm) generally falling in June.

The Clarion-Nicollet-Webster soil association, which is the only major soil group in Test Site 3, was derived from glacial till. About 75 percent of the area has level to gently sloping topography and is well suited to intensive production of corn, soybeans, and alfalfa.

This test site has about 1,499,600 hectares of farm land and typically grows 607,300 hectares of corn (approximately 96% for grain); 477,100 hectares of soybeans; and 54,500 hectares of alfalfa and hay.

Three major kinds of parent materials (loess, glacial till, and alluvium) are found in Test Site 4. Loess (wind-blown material) from the Missouri flood plains is thickest near the Missouri River and thins and increases in clay content in a southeasterly direction. Marshall and Monona-Ida-Hamburg soil associations which occupy the central three-fourths of this district were formed from deep loess under grass vegetation. These soils are generally well-drained and have high proportions of their area used for cultivated crops. The Clarion-Nicollet-Webster soil association, which is a continuation of the predominant soil of the third test site, is the major soil in Sac County. These soils are well suited to intensive production of corn, soybeans, and alfalfa. A third major group of soils which developed primarily from alluvial materials on the nearly level flood plains of the Missouri River are the Luton-Onawa-Salix association. These soils are found primarily along the Missouri River in Woodbury, Monona, and Harrison counties and are farmed for corn, soybeans, and wheat.

High proportions of Test Site 4 are used for cultivated crops, particularly corn and soybeans. Of the 1,385,100 hectares of farm land in this district, 634,100 hectares of corn are planted annually and approximately 90 percent of this corn is harvested for grain. An additional 233,700 hectares of soybeans are typically planted. The proportions of corn and soybeans vary from year to year depending on market conditions and prices.

#### Sample Allocation.

In general, two segments per county were allocated. In the case of unusually large or small counties, three segments or one segment might be allocated. All counties indicated in Figure 12 received segments. Table 12 lists the number of segments allocated to each county.

Table 12. Allocation of sample segments to counties in each of the four high density test sites.

Test Sites	State	County	No. of Segments
1	Indiana	Adams	2
		Allen	2
		Blackford	2
		Delaware	2
		Henry	2
		Jay	2
		Madison	2
		Randolph	2
		Wayne	2
		Wells	2
2	Indiana	Benton	2
		Jasper	2
		Newton	2
		Tippecanoe	2
		Warren	2
	Illinois	Champaign	3
		Ford	1
		Iroquois	3
		Kankakee	2
		Vermilion	3
3	Iowa	Calhoun	2
		Emmet	2
		Hamilton	2
		Hancock	2
		Humboldt	2
		Kossuth	2
		Palo Alto	2
		Pocahontas	2
		Webster	2
		Wright	2
4	Iowa	Crawford	2
		Harrison	2
		Ida	2
		Monona	2
		Pottawatomie	3
		Sac	2
		Shelby	2
		Woodbury	3

### Sample Location.

The method used for sample selection was the same as described for the low density samples. More segments were located than were allocated to permit for loss of some segments in nonagricultural areas. Locations of the sample segments by latitude and longitude coordinates can be found in Table 13.

### 7. Summary and Conclusions

A stratification was performed and sample segments were selected for an initial investigation of Multicrop problems. The effort was to support:

- Development and evaluation of procedures for using LACIE and other technologies for the classification of corn and soybeans.
- Identification of factors likely to affect classification performance.
- Evaluation of problems encountered and techniques which are applicable to the crop estimation problem in foreign countries as well.

The two types of samples, low density and high density, supporting these requirements were selected as a research data set for an initial evaluation of technical issues and should not be used in an aggregation scheme. In summary, looking at the geographic location of the strata, the system appears to be logical and the various strata seem to represent different conditions. This result is supportive not only of the variables and the methodology employed in the stratification, but also of the validity of the data sets employed.

Table 13. Locations of the high density sample segments by latitude and longitude coordinates of the segment centers.

Test Site	State	County	Latitude/Longitude
1	Indiana	Adams	40.785/84.880
		Adams	40.620/85.016
		Allen	40.956/85.273
		Allen	40.952/84.877
		Blackford	40.541/85.412
		Blackford	40.457/85.413
		Delaware	40.290/85.285
		Delaware	40.123/85.549
		Henry	39.789/85.424
		Henry	40.039/85.419
		Jay	40.370/85.022
		Jay	40.451/84.889
		Madison	40.128/85.810
		Madison	40.295/85.808
		Randolph	40.038/85.159
		Randolph	40.036/84.899
		Wayne	39.785/84.904
		Wayne	39.954/85.161
		Wells	40.789/85.276
Wells	40.650/85.230		
2	Indiana	Benton	40.627/87.382
		Benton	40.520/87.210
		Jasper	40.879/86.990
		Jasper	40.963/87.122
		Newton	41.125/87.521
		Newton	40.794/87.384
		Tippecanoe	40.515/87.027
		Tippecanoe	40.335/86.835
		Warren	40.378/87.117
		Warren	40.293/87.378
	Illinois	Champaign	40.172/88.307
		Champaign	40.339/88.435
		Champaign	40.310/88.100
		Ford	40.588/88.300
		Iroquois	40.919/88.030
		Iroquois	40.752/88.034
		Iroquois	40.831/87.768
		Kankakee	41.248/87.757
		Kankakee	41.086/88.026
Vermilion	40.078/87.657		
Vermilion	40.415/87.910		
Vermilion	40.330/87.650		

Table 13 (con't)

---

Test Site	State	County	Latitude/Longitude
3	Iowa	Calhoun	42.294/94.838
		Calhoun	42.380/94.569
		Emmet	43.464/94.725
		Emmet	43.298/94.585
		Hamilton	42.219/93.489
		Hamilton	42.300/93.893
		Hancock	43.052/93.625
		Hancock	43.135/93.762
		Humboldt	42.801/94.036
		Humboldt	42.717/94.303
		Kossuth	42.966/94.301
		Kossuth	43.135/94.172
		Palo Alto	42.963/94.852
		Palo Alto	43.127/94.855
		Pocahontas	42.713/94.711
		Pocahontas	42.794/94.848
		Webster	42.384/94.164
		Webster	42.549/94.166
		Wright	42.886/93.897
		Wright	42.217/93.876
4	Iowa	Crawford	41.948/95.635
		Crawford	41.952/95.367
		Harrison	41.615/95.624
		Harrison	41.778/95.763
		Ida	42.454/95.382
		Ida	42.530/95.655
		Monona	41.941/96.037
		Monona	42.113/95.775
		Pottawatomie	41.285/95.348
		Pottawatomie	41.446/95.619
		Pottawatomie	41.362/95.749
		Sac	42.205/95.374
		Sac	42.457/95.111
		Shelby	41.699/95.493
		Shelby	41.622/95.224
		Woodbury	42.358/96.054
Woodbury	42.216/95.784		
Woodbury	42.480/95.870		

---



APPENDIX

#### Appendix. Problems Encountered with County Estimates Data

Numerous difficulties were encountered with the county estimates data. The original tape which was transmitted from NASA/JSC was in ASCII format on an 800 BPI tape.

There were some unreadable characters on the tape, indicating that the original data tape may have been bad. The problems were found in the first few columns of a record, so that "educated guesses" could be made to fill in the missing information. The missing information was sometimes restricted to the first five columns which were constant throughout the entire data set. If state or county codes were missing, these could be determined by examining the placement of the card in the data deck. For all bad data lines, the missing information was overlaid, but the first five columns (containing a constant code which was irrelevant to the study) were left bad so that these lines could be located again if necessary.

There were also some codes encountered which were not documented. A visit with Bob Cole of the Indiana USDA/SRS office helped identify an appropriate course of action.

The first column of each record was supposed to indicate the card number and should have been "2" for all data on the tape. One record was encountered, however, which had "3" in the first column. As it was learned that card three did not exist, this was determined to be a key-punching error and was changed on the data file.

Table A-1 lists crops and their codes for the data set. Nonexistent commodity codes were encountered in the data files. Some of the unusual codes might have been misspunched or might have been specific to a state; an example of this type is the code 17163 (for class and crop code). Class code 33 was not included on the list in Table A-1, but the Indiana office of the USDA/SRS was able to inform us that this class code represented

PRECEDING PAGE BLANK NOT FILMED

44

Table A-1. Commodity Codes.

Class	Crop Code	Utili- zation	Crop Name
10	119	9	Winter Wheat
10	129	9	Durum Wheat
10	139	9	Other Spring Wheat
10	199	9	Wheat, All
10	499	9	Rye, All
10	619	9	Rice, All
11	199	1	Corn for Grain
11	199	2	Corn for Silage
11	299	9	Oats, All
11	399	9	Barley, All
11	499	9	Sorghum, All
12	129	9	Cotton, All (Neither Ginning Status nor Staple Type Specied)
12	121	9	Cotton, Upland
12	122	9	Cotton, American Pima
			<u>Tobacco:</u>
14	111	1	Flue-cured, type 11
14	111	2	Flue-cured, type 12
14	111	3	Flue-cured, type 13
14	111	4	Flue-cared, type 14
14	122	1	Fire-cured, type 21
14	122	2	Fire-cured, type 22
14	122	3	Fire-cured, type 23
14	133	1	Air-cured, type 31
14	133	2	Air-cured, type 32
14	133	5	Air-cured, type 35
14	133	6	Air-cured, type 36
14	133	7	Air-cured, type 37
14	244	1	Cigar-filler, type 41
14	255	1	Cigar-binder, type 51
14	255	2	Cigar-binder, type 52
14	255	4	Cigar-binder, type 54
14	255	5	Cigar-binder, type 55
15	299	9	Flaxseed
15	399	1	Peanuts
15	499	1	Soybeans
16	171	1	Dry Edible Beans - Pea (Navy)
16	171	2	- Great Northern
16	171	4	- Flat Small White
16	171	6	- Pinto
16	171	7	- Red Kidney
16	172	1	- Pink
16	172	2	- Small Red
16	199	9	Dry Beans (All Mich.)
16	319	9	Dry Peas - Smooth Green Kinds
16	329	9	- Yellow and White Kinds
83	161	8	Wrinkled Peas for Seed
16	599	9	Lentils
16	819	9	Austrian Winter Peas
36	129	9	Green Peas for Processing
37	829	9	Tomatoes for Processing
83	104	2	Bush Garden Seed Beans (Idaho)

miscellaneous vegetables. Another problem was class codes which matched the list given, but whose corresponding crop code or utilization code did not exist. The code "14558", for example, does not exist, but all of class 14 is tobacco so this observation was included there. The utilization "0" is not used for 10129 (durum wheat) but was included in that crop type anyway. There were approximately 40 more cases which were handled in a similar fashion.

Duplicate cards were also encountered in the winter wheat, corn, barley, and miscellaneous crops data files. There were several different varieties of duplicates. Some cards were exact duplicates, a situation which had a straightforward solution. Some cards were encountered containing different estimates of a crop for the same county and the same year, but which were punched in different years. In this case, the most recently entered information was selected to be correct. Some duplicate cards had a third type of problem: yields differed by a factor of ten while the rest of the information was identical. In this situation, the card was selected for which acres times yield was equal to production.

There were many zeros for acreage, yield, and production in the data files. By looking at the values for a given crop in a given county over the five year period, it was determined that a zero might represent two situations: either no acreage of that crop was grown in that county or the true data value was missing. Missing values could possibly have been determined by consulting state crop production publications; time constraints for this project, however, did not permit this type of verification. Years with zero values were excluded from consideration in computation of crop averages. If the data were indeed missing, this approach yielded a much more realistic estimate. If the data were truly zero, a good estimate would be obtained by averaging the other non-zero years which would be small numbers.

Additional steps in data verification were attempted by summing individual estimates to obtain a total for a given crop, crop reporting

district, or state. These methods were abandoned as a data verification aid again due to resource considerations. As a rule, summing did not work. Some examples are given as follows.

Theoretically, the "all wheat" data file should be the sum of the winter wheat, durum wheat, and other spring wheat files. This was supposed to be true according to Bob Cole of the Indiana USDA/SRS office, but was found to be not necessarily true in the data. Sometimes all wheat was larger than the sum of the three component files and sometimes the sum was larger. Occasionally, the numbers were about equal. Some counties reported total wheat, but did not divide it down into its components, while other counties appeared to do the reverse.

Finally, crop reporting district and state area and production estimates of a crop should be the sum of estimates for the counties comprising them. This check also failed frequently, a possible result of missing data or a mixture of preliminary and final estimates.