

Reprinted from

**Symposium on  
Machine Processing of  
Remotely Sensed Data**

**June 21 - 23, 1977**

The Laboratory for Applications of  
Remote Sensing

Purdue University  
West Lafayette  
Indiana

IEEE Catalog No.  
77CH1218-7 MPRSD

Copyright © 1977 IEEE  
The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to [pubs-permissions@ieee.org](mailto:pubs-permissions@ieee.org).

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

## MONITORING EARTH ALBEDO FROM LANDSAT

RICHARD K. KIANG AND STEPHEN G. UNGAR

NASA Institute for Space Studies  
Goddard Space Flight Center  
New York, New York 10025

A technique has been developed for extracting ground albedo from LANDSAT data. Initial application to Central Kansas indicates a trend of increasing ground albedo superimposed on seasonal variations. The measured albedo apparently is associated with drought. The results suggest that LANDSAT data can be used as an input to global climatic prediction models which depend on ground albedo information.

The technique for obtaining the albedos uses a doubling radiative transfer model with scaling approximation. Reflectivities are extrapolated beyond the LANDSAT MSS spectral region based on categorization of the scene into ground classes using the original LANDSAT data. The extrapolation factors have been derived from empirical signatures. The proportions of ground classes are determined from supervised classification when more than one class is present. A cloud detection algorithm is used to remove the cloud contaminations from LANDSAT measurements.

The shortwave radiation measurements with low spatial resolution from weather satellites such as Nimbus III and Nimbus 6 had been used to calculate earth albedo in synoptic scale. Because of LANDSAT's much higher spatial resolution, its measurements are particularly useful in calculating the total area of each climatic system, monitoring natural and anthropogenic impact on the earth's surface, and preparing a detailed albedo atlas.