

Reprinted from

Symposium on

Machine Processing of

Remotely Sensed Data

June 27 - 29, 1979

The Laboratory for Applications of
Remote Sensing

Purdue University
West Lafayette
Indiana 47907 USA

IEEE Catalog No.
79CH1430-8 MPRSD

Copyright © 1979 IEEE
The Institute of Electrical and Electronics Engineers, Inc.

Copyright © 2004 IEEE. This material is provided with permission of the IEEE. Such permission of the IEEE does not in any way imply IEEE endorsement of any of the products or services of the Purdue Research Foundation/University. Internal or personal use of this material is permitted. However, permission to reprint/republish this material for advertising or promotional purposes or for creating new collective works for resale or redistribution must be obtained from the IEEE by writing to pubs-permissions@ieee.org.

By choosing to view this document, you agree to all provisions of the copyright laws protecting it.

APPLICATION OF A PRINCIPAL COMPONENTS ANALYSIS ON LANDSAT MULTISPECTRAL DATA FOR STUDIES ON VEGETATION COVER UNDER DESERT CONDITIONS

JELLE U. HIELKEMA

Food and Agriculture Organization of the United Nations

Present day available satellite remote sensing technology offers the possibility of obtaining rapid and comprehensive information about the earth's natural resource base at relatively low cost. Especially for conventional data sparse and often inaccessible remote areas these new intelligence gathering techniques are of considerable interest. The present limited spatial and spectral resolution of the Landsat multispectral scanners poses a problem to the effective recording of low coverages of vegetal growth in desert environments. The spectral response from the vegetation is generally contaminated with the reflected radiation from the abiotic background.

This paper discusses the application of a principal components transformation on Landsat multispectral data for enhancement of small areas with low coverages of ephemeral vegetation in an arid environment.

In arid areas the four Landsat spectral bands exhibit a very high degree of correlation. This characteristic, combined with the basic differences between the spectral properties of vegetation and bare surface materials, makes that a principal components transformation on Landsat multispectral data can be successfully used as an image enhancement technique. By application of a principal components transformation, very low coverages of desert vegetation can be detected by visual analysis; such vegetation coverages cannot be extracted from false colour composite imagery constructed from original 4, 5, 7 channel data.

Utilization of a principal components analysis as a data reduction technique for multispectral data of arid areas is also demonstrated.